

Available online at www.sciencedirect.com



JOURNAL OF GENETICS AND GENOMICS

J. Genet. Genomics 36 (2009) 325-334

www.jgenetgenomics.org

Liver development in zebrafish (Danio rerio)

Ting Tao, Jinrong Peng^{*}

College of Animal Science, Zhejiang University, Hangzhou 310029, China

Received for publication 16 February 2009; revised 17 April 2009; accepted 17 April 2009

Abstract

Liver is one of the largest internal organs in the body and its importance for metabolism, detoxification and homeostasis has been well established. In this review, we summarized recent progresses in studying liver initiation and development during embryogenesis using zebrafish as a model system. We mainly focused on topics related to the specification of hepatoblasts from endoderm, the formation and growth of liver bud, the differentiation of hepatocytes and bile duct cells from hepatoblasts, and finally the role of mesodermal signals in controlling liver development in zebrafish.

Keywords: zebrafish; model organism; liver development; endoderm

Introduction

Liver is an essential organ in the body and performs a number of vital activities including metabolism, detoxification and homeostasis. Hepatocytes make up the majority of a liver and exert most of the liver's functions, such as metabolism of a wide range of endogenous and exogenous substances, synthesis of blood protein and clotting factors, storage of glycogen, amino acids, fat, and iron, as well as secretion of bile (Blouin et al., 1977; Kawamoto et al., 1996; Pack et al., 1996; Field et al., 2003). Liver diseases including hepatitis B, cirrhosis and hepatocellular carcinoma (HCC) are worldwide health problems and cause high death toll annually, making great threaten to human health and social development (Dienstag, 2008). Therefore, it is important to illustrate liver development and its cellular and molecular mechanism. The knowledge learned will not only generate new approaches for early monitoring, prevention and therapy of the liver diseases but also help to guide in vitro culture and transplantation of the liver. In

this review, we briefly summarized the major progresses in zebrafish (Danio rerio) in the following aspects of liver development: the origination and specification of hepatoblasts within endoderm, the formation and growth of liver bud, the differentiation of hepatocytes and biliary cells from hepatoblasts and the roles of mesodermal signals during the liver development.

Zebrafish as an excellent model system for studying liver development

Zebrafish belongs to teleosts whose liver has distinguishable histological characteristics compared with that in the mammalian. Its portal veins, hepatic arteries and large biliary ducts are distributed stochastically within the hepatic parenchyma but are not grouped in portal tracts as in the mammalian. Besides, hepatocytes are arranged as tubules that enclose small bile ducts rather than as bilayered hepatocyte plates appeared in the mammalian. The intrahepatic bile ducts are derived from the bile cannaliculi and form a network of biliary channels; the bile is then col-

^{*} Corresponding author. Tel & Fax: +86-571-8697 1513. E-mail address: pengjr@zju.edu.cn

lected in the gallbladder through large ducts and an extrahepatic biliary system (Pack et al., 1996; Lorent et al., 2004).

In mammals, liver organogenesis begins with the establishment of a population of hepatic precursor cells within the ventral foregut endoderm, followed by the specification of definitive hepatoblasts. The hepatoblasts delaminate from the epithelial layer to form a discrete liver bud, and undergo rapid proliferation to increase the size of the bud. Finally, the hepatoblasts differentiate into functional hepatocytes and biliary duct cells (Zaret, 2002; Duncan, 2003). Although the process of liver initiation and development at the anatomic level is well established, its molecular and cellular mechanisms are relatively understudied through genetic approaches, and only limited information has been obtained up to now. The study of liver development is mainly constrained by the model animals used such as mouse and chicken, which cannot be used for large-scale genetic analysis. Zebrafish, on the other hand, has become a popular vertebrate model during last decade because both genetic and experimental embryological methods can be easily applied to this animal (Grunwald and Eisen, 2002). Zebrafish is a small tropical freshwater fish native to South Asia and is a common aquarium fish around the world. In 1981, Streisinger and his colleagues published a landmark paper describing the genetic procedures for producing clones of homozygous diploid zebrafish (Streisinger et al., 1981). At the Cold Spring Conference in 1994, zebrafish was formally established as a model animal for the vertebrate development study (Concordet and Ingham, 1994; Kahn, 1994; Mullins et al., 1994; Solnicakrezel et al., 1994). In 1996, an entire issue of 'Development' journal reported two independent large-scale mutagenesis screens for phenotypic mutants and the analysis of some of the obtained mutants in zebrafish (Driever et al., 1996; Haffter et al., 1996). With more and more exciting findings published in a wide range of journals, zebrafish has shown its great potential in life science. Now it has become an ideal model organism for the study of vertebrate development and disease, functional genomics, organ function, behavior, toxicology and drug discovery. This is mainly attributed to some of its advantages, including its easy and economical maintenance, short generation time, external fertilization, large numbers of rapidly developing embryos produced per mating, and the external development of transparent embryos (Streisinger et al., 1981; Grunwald and Eisen, 2002). Given that fundamental developmental programs are well shared among vertebrate animals, the studies on zebrafish

should naturally contribute to a better understanding of the molecular and genetic mechanisms of development underlying other classes of vertebrates including human.

In addition to the above mentioned characteristics, zebrafish has some unique advantages for studying liver development. In mammals, the embryonic liver is an early hematopoietic organ, therefore, mutations affecting liver or blood development often cause anemia and even early lethality during embryogenesis that will complicate the study of liver development in the mammalian system (Reimold et al., 2000). Furthermore, mammalian embryogenesis occurs intrauterinely which makes the embryonic liver inaccessible for the study of the processes of liver development via a direct genetic approach. On the other hand, the blood (primitive) in zebrafish forms in the ICM (intermediate cell mass) first and subsequently (definitive) in the PBI (posterior blood land) and lastly kidney, not in the liver (Detrich et al., 1995; Thisse and Zon, 2002; Jin et al., 2009). Meanwhile, since zebrafish obtains its nutrient mainly from yolk during embryogenesis, zebrafish can survive and go on to develop relatively normally for a few days without the cardiovascular system (Stainier, 2001). These advantages allow the studies on liver development and disease in zebrsfish even the mutations affecting blood development. Last but not least, the availability of a number of transgenic fish lines harboring the reporter genes gfp and/or rfp driven by a variety of endoderm-specific promoters or the heat-shock promoter not only allow researchers to follow the developmental processes of the liver in vivo but also greatly facilitate mutant screening based on the expression of the reporter genes (Table 1).

The origination and specification of hepatoblast

The endoderm gives rise to the hepatoblast. Fate mapping experiments have revealed that, at early gastrula stage (6 hpf, hours post-fertilization), endoderm cells located in the ventral part tend to differentiate to liver bud while endoderm cells situated on the dorsal side are apt to give rise to pancreas. These liver-only progenitors are located considerably further from the dorsal organizer than the pancreas-only progenitors. Clones that contribute to both liver and pancreas are prone to lie in more intermediate positions (Ward et al., 2007).

Currently, there are two different hypotheses regarding the initiation of liver progenitor cells in zebrafish. Based on the fact that the liver specific marker ceruloplasmin (cp) can be detected in the dorsal endoderm at 16 hpf which is

Table 1
Transgenic reporter fish used in the studies of liver development in zebrafish

a 1		
Transgenicline	Tissue expression	References
Tg(gutGFP) ⁵⁵⁸⁴	Express the gfp gene in the entire alimentary canal and endodermal component of all accessory organs	Field et al., 2003
Tg(sox17:GFP) ^{s870} ; Tg(sox17:DsRed) ^{s803}	Express the gfp and rfp genes in the sox17 positive endoderm cells, respectively	Chung and Stainier, 2008
Tg(Ifabp:EGFP); Tg(Ifabp:dsRed)	Express the gfp and rfp genes in the liver, respectively	Her et al., 2003; Shin et al., 2007
Tg(hsp701:dnBmpr-GFP) ^{w30}	Express a dominant negative form of the Xenopus Bmp receptor type 1a fused to GFP after heat shock	Pyati et al., 2005, 2006
Tg(hsp70:noggin3)	Express Noggin3 after heat shock	Chocron et al., 2007
Tg(hsp701:bmp2b) ^{f13}	Express Bmp2b after heat shock	Chocron et al., 2007
Tg(hsp70:alk8)	Express alk8 (a type I Bmp receptor) after heat shock	Shin et al., 2007
Tg(hsp70l:dnfgfr1-EGFP)	Express a dominant negative form of Fgfr1 fused to GFP after heat shock	Lee et al., 2005b
Tg(ptf1a:GFP) ^{ih1}	Express GFP in the developing ventral pancreas	Godinho et al., 2005; Chung et al., 2008

prior to the liver morphogenesis, Korzh et al. (2001) hypothesized that liver progenitors might be differentiated before the initiation of alimentary canal morphogenesis and the liver bud was formed later by migration and aggregation of these liver progenitor cells. Similar hypothesis is also applied to the pancreas initiation because the pancreas marker pdx1 is first detected at the 10-somite stage (~14 hpf) which is way ahead of the pancreas morphogenesis (Biemar et al., 2001; Wallace and Pack, 2003). However, so far there is no concrete evidence to demonstrate that the early cp positive cells do contribute to the formation of liver bud at a later stage. As for the pdx1 positive cells, it is likely that they mainly contribute to the endocrine pancreatic cells but not to the exocrine pancreas (Field et al., 2003). The second hypothesis is based on the fact that the expression of two transcription factors hhex and prox1, two key hepatoblast markers, are detectable in the endoderm region that later gives rise to liver bud, suggesting that liver progenitor cells are differentiated from the endoderm cells to form in situ liver bud after the formation of endoderm rod. The latter hypothesis is supported by data from anatomic studies (Field et al., 2003; Wallace and Pack, 2003) and by studies on mutants related to liver development (Ober et al., 2006; Shin et al., 2007; Huang et al., 2008).

Liver bud formation and growth

Liver is an accessory organ of the foregut. Based on the results observed in the GutGFP transgenic zebrafish Tg(gutGFP)⁵⁵⁸⁴ (expressing the gfp reporter gene in the

entire alimentary canal and endodermal component of all accessory organs), Field et al. (2003) proposed that the liver morphogenesis process can be arbitrarily divided into two phases: budding and growth. The budding phase occurs from 24 to 50 hpf and can be further divided into three stages. At the budding stage I, endoderm cells caudal to the pharyngeal region aggregate to form endoderm rod at 24 hpf. At 28 hpf, the endoderm rod segment under the first somite starts to thicken, which marks the beginning of liver morphogenesis. From 28 hpf (budding stage II), the anterior thickening region, which refers to liver primordium, increases in size and bends to the left side (a process termed as gut-looping) with respect to the middle line (Horne-Badovinac et al., 2003), and then covers the outer curvature of intestinal bulb by 30 hpf. The formation of a furrow between the liver bud and the adjacent esophagus marks the beginning of last phase of liver budding at approximately 34 hpf. Along with the expansion of the furrow, the hepatic duct is shaped to connect liver and intestine at the end of budding stage III. The liver now appears to locate between the duct of Cuvier, anteriorly, and the mid-level of fin bud, posteriorly (Field et al., 2003) (Fig. 1).

At the subsequent growth stage, the liver undergoes dramatic changes in its size, shape and placement because of rapid cell proliferation. Hepatoblasts also start to differentiate into functional hepatocytes and bile duct cells. By 96 hpf, the liver extends across the midline ventral to esophagus to form the second liver lobe (right lobe). It touches the pericardial cavity anteriorly and overlaps with the anterior portion of residual yolk (Field et al., 2003) (Fig. 1). Cellular and molecular control of liver development in zebrafish

In merely less than a decade, the power of using zebrafish



Fig. 1. Dorsal view of liver development in zebrafish. The foxa3 probe was used in whole-mount in situ hybridization (WISH) for embryos at 30, 34 and 48 hpf as shown (black arrow: liver; red arrow: pancreas). The Ifabp probe was used in WISH for embryos at 72, 96 and 120 hpf. A, anterior; P, posterior.

to study liver development has been demonstrated by a number of high-quality publications. In the following, we summarized the main achievements of these studies (Fig. 2).

Signaling molecules (Bmp, Fgf, Wnt, Shh and RA)

Early in 1975, Douarin found that hepatocytes differentiated from endoderm required interactions with adjacent precardiac mesoderm in chick (Douarin, 1975), and the same result was confirmed in mouse (Gualdi et al., 1996). It was ascribed to the signaling molecules such as fibroblast growth factors (Fgfs) from the cardiac mesoderm (CM) (Jung et al., 1999) and bone morphogenetic proteins (Bmps) from the septum transversum mesenchyme (STM) (Rossi et al., 2001). During the budding stage in zebrafish, liver cells are observed adjacent to lateral plate mesoderm (LPM) cells at the dorsal face and to yolk ball at the ventral face directly (Horne-Badovinac et al., 2003). Therefore, the question arises whether zebrafish LPM cells express signaling molecules functioning to regulate liver development similar to Fgf and Bmp signaling in mouse and chick. The essential role of the mesoderm tissues for liver develop-f2t(m)ules e Ting Tao et al. /

type. Therefore, as found in mouse (Zhao et al., 2005), Gata4 and Gata6 have redundant function during early stage of liver development and are both essential for liver growth in zebrafish (Holtzinger and Evans, 2005).

Retinol binding protein 4 (rbp4) is not only expressed in the adult zebrafish liver, but also detected in the yolk syncytial layer (YSL). rbp4 is negatively mediated by Nodal and Hedgehog (Hh) signaling, but positively regulated by RA signal. It regulates liver budding and may be also required for migration of liver primordia. rbp4 morphants show two liver buds which may be due to improper migration of liver primordia. Since rbp4 also regulates the activity of fibronectin I (fnI), fnI may act on liver primordia migration directly (Li et al., 2007). Knockdown of vegfc in zebrafish embryos also results in multiple liver and pancreatic buds; however, the expressions of both anterior liver bud and posterior pancreatic bud markers are unaffected, indicating that the anteroposterior patterning and organ differentiation of the gut endoderm is normal (Ober et al., 2004). Studies on the zebrafish cloche (clo) mutant show that endothelial cells do not appear to be essential for liver budding (Field et al., 2003), which is different from what have been observed in mouse (Matsumoto et al., 2001).

Def (digestive-organ-expansion-factor), a novel nuclearlocalized protein, is required for digestive organ expansion but not organ initiation and cell differentiation. Loss-offunction of def selectively up-regulates the expression of a newly identified p53 isoform Δ 113p53 (human counterpart Δ 133p53) that leads to up-regulation of cell-cycle-related genes but not apoptotic-related genes to arrest the expansion growth of major digestive organs, including liver (Chen et al., 2005, 2009). Two other factors, nil per os (npo) encoding an RNA-binding protein and leg1 (liver-enriched gene 1), have also been shown to play a crucial role in the expansion growth of embryonic liver in zebrafish (Mayer and Fishman, 2003; Cheng et al., 2006).

Pescadillo (Pes) is highly conserved from yeast to human and can be detected firstly at 48 hpf in the liver primordium, followed by the reduced RNA level at 72 hpf. During normal development between 72 hpf and 144 hpf, the liver extends over the yolk surface and this process is associated with rapid consumption of the yolk. On the other hand, in the pes mutant (generated via retroviral vector insertional mutagenesis method) the liver expansion is arrested beginning at 72 hpf, the mutant yolk is not consumed and the mutant fish dies at 144 hpf (Allende et al., 1996).

Expression of the ubiquitin-like protein containing PHD

and ring finger domains-1 (uhrf1) gene is enriched in the liver bud and fully developed liver but is rare in the adult zebrafish liver. Analysis of the zebrafish hi272 line, which bears an insertion in uhrf1, reveals that uhrf1 regulates liver apoptosis and proliferation but has no visual effects on hepatocytes histological manifestation. In the uhrf1 mutant, the liver is shaped as a ball whereas the right form should be crescent-shaped in wild type at 120 hpf. In addition, uhrf1 also functions in zebrafish liver regeneration (Sadler et al., 2007).

Some other factors have also been identified to play roles in the liver development. Rai et al. found that DNA methyltransferase 2 (Dnmt2) (Rai et al., 2007) but not Dnmt1 (Rai et al., 2006) was required for liver development in zebrafish. Dnmt2 is a maternal protein and its specific expression in liver is only detectable at 72 hpf. Since the expression of late but not early liver differentiation markers are impaired in the dnmt2 morphants, Dnmt2 may affect the late stage of hepatocyte differentiation through cytoplasmic RNA methylation (Rai et al., 2007). Ftz-f1 belongs to the nuclear receptor type of transcription factor and is expressed in different forms. The IIA form of Ftz-f1 is the major transcript during zebrafish early liver morphogenesis, indicating its potential role in liver development (Lin et al., 2000). Although no direct evidence is obtained up to now, the function of Ftz-f1 homologs in human (Li et al., 1998) and rat (Galarneau et al., 1996) has indicated this notion. HNF-1 is a vertebrate transcription factor that contains a divergent homeodomain. In the zebrafish hnf1 mutant (vhnf1^{hi548} allele), liver is poorly formed with undifferentiated hepatocytes at 72 hpf (Sun and Hopkins, 2001). The liver marker gene hhex, which is also essential for liver formation (Wallace et al., 2001), expresses at a low level in the vhnf1^{hi548} mutant, indicating its role downstream of vHnf1. It is proposed that vHnf1 controls liver development through regulating the balance of the pdx1-shh network (Sun and Hopkins, 2001). Recently, Cheng et al. (2006) reported their analysis of promoter regions of 51 liver-enriched genes by searching putative binding sites for Hnf1, Hnf3, Hnf4 and Hnf6, and revealed that these four liver-enriched transcription factors form a network to control the expression of liver-specific or liver-enriched genes in the liver. Interestingly, chromatin remodeling factors histone deacetylase 1 (Hdac1) (Noel et al., 2008) and Hdac3 (Farooq et al., 2008) have also been shown to play important but relatively distinct roles during liver specification and development in zebrafish. It seems that Hdac3 acts through inhibiting its unique target growth differentiation factor 11 (Gdf11), a member of the

331

transforming growth factor beta family, to promote liver development (Farooq et al., 2008).

Genes essential for differentiation of hepatocytes and bile duct cells

Onecut transcription factors (Oc) play important roles during zebrafish liver and pancreas development. onecut1 (oc1, mammalian ortholog hnf6) and its downstream gene vhnf1 are both essential for the biliary system development. Knockdown or overexpression of either gene perturbs the development of intrahepatic but not extrahepatic bile ducts (Matthews et al., 2004). As the third member of onecut gene family and also a functional ortholog of mammalian hnf6, onecut3 (oc3) regulates the earliest stages of zebrafish biliary development. Expression of oc3 is obviously reduced in oc1-deficient embryos, and vice versa. Therefore, members of the Onecut family seem to interact with each other to regulate biliary development (Matthews et al., 2004).

In addition, jagged 1, 2, 3 and notch 1a, notch 1b, notch 2, and notch 5 are each expressed in liver at 48 hpf and 72 hpf, exactly the time when bile ducts form. Absence of jagged or notch genes affects a series of zebrafish tissues including biliary system development, causing a phenotype comparable with the human Alagille Syndrome (AGS) (Lorent et al., 2004). Disruption of the biliary and pancreatic ductular system also appeared in hhex morphant (Wallace et al., 2001).

Genes essential for hepatopancreatic duct development

During morphogenesis of liver and exocrine pancreas, liver bud and exocrine pancreatic bud are linked by the hepatopancreatic duct. However, the nature and function of the hepatopancreatic duct have been studied only recently. In the zebrafish fgf10 mutant, the hepatopancreatic duct epithelium becomes malformation and the contiguous cells fate is disordered (Dong et al., 2007). Based on the detailed molecular and cellular study, it is established that Fgf10 secreted from the adjacent mesenchyme functions not only to refine the boundaries between the hepatopancreatic duct and organs, but also to prevent the differentiation of the proximal hepatic and pancreatic cells into each other (Dong et al., 2007). Interestingly, in addition to its role in liver organogenesis, a distinct role of histone deacetylase 1 (Hdac1) in extrahepatopancreatic duct morphogenesis has also been identified (Noel et al., 2008).

Zebrafish as a model for liver disease study

In addition to its contribution to the study of liver development, zebrafish is increasingly being used to study liver diseases. Lam et al. (2006) generated liver tumors in zebrafish by treating the fish with carcinogens and then obtained the gene set for liver tumors via microarray hybridization. Next, they cross-compared the zebrafish liver tumor gene set with the gene sets obtained from four cancer types (liver, gastric, prostate and lung) in human. Astonishingly, they found that the zebrafish liver tumor gene set intersects most with the gene set from human liver tumors and this lays the foundation to establish zebrafish as a model system to study liver tumorogenesis (Lam et al., 2006). The fact that -hexachlorocyclohexane, thioacetamide and alcohol each can induce hepatic steatosis in zebrafish successfully convinces the scientific community that zebrafish is a good model system to study steatohepatitis (Braunbeck et al., 1990; Amali et al., 2006; Passeri et al., 2009). More importantly, zebrafish serves as a genetic model system for studying the molecular mechanism behind hepatic steatosis. For example, it has been shown that mutation in a novel gene foie gras leads to hepatomegaly likely due to enlarged lipid-filled hepatocytes in the mutant liver. Detailed study of the gonzo mutant (mutated in the membrane-bound transcription factor protease 1 gene) showed that alcohol induced hepatic steatosis was largely through activation of the pathway mediated by the sterol response element binding protein (SREBP) transcription factors (Passeri et al., 2009). On the other hand, mutation in the S-adenosylhomocysteine hydrolase (ahcy) gene increases that expression to cause the hepatic steatosis phenotype in the ducttrip (dtp) mutant (Matthews et al., 2009). In a screen for zebrafish hepatomegaly mutants, Sadler et al. (2005) found that mutations in a class C vacuolar sorting protein gene vps18 and a tumor suppressor gene nf2 both conferred hepatomeglay due to defects in the biliary system development.

Conclusion

In recent years, zebrafish has shown a full potential in the studies of vertebrate liver development. The combination of forward and reverse genetics studies, and classical genetics analysis coupled with direct embryos observation and manipulation, as well as its particular suitability for the study of liver organogenesis, will no doubt strengthen our understanding of the cellular and molecular mechanism of liver development.

Liver formation is a complicated process. Thus, there must be multiple genes and signaling molecules involved in this process. Up to now, we have known that Fgf, Bmp, Wnt and RA signaling pathways are involved in the initiation and differentiation of hepatocyte in zebrafish, but the precise relationship among them remains unclear. Furthermore, only a handful of factors had been identified to affect liver budding and growth. For example, hhex acts to control the initiation and budding of liver progenitors. prox1 is required for the outgrowth of liver bud and migration of hepatoblasts whilst members of gata family, as pan-endodermal factors, are crucial for liver development. However, these few factors and pathways are not sufficient for understanding the whole process of liver initiation and development. Therefore, while further studies of known genes and existing mutants will provide us with new information for the understanding of liver organogenesis, more liver-defective mutants need to be identified for illustrating mechanisms that are behind the mutant phenotype observed. The identification of such mutants would lead to revealing the genes that are crucial for liver organogenesis. Though the method for target gene knockout in zebrafish is not sophisticated yet at present, we believe breakthrough will be made in the near future. Apparently, knowledge gained from studying liver development will help to guide the fields such as liver transplantation, gene therapy and tissue engineering, and to benefit human being as a whole.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 30825025).

References

- Allende, M.L., Amsterdam, A., Becker, T., Kawakami, K., Gaiano, N., and Hopkins, N. (1996). Insertional mutagenesis in zebrafish identifies two novel genes, pescadillo and dead eye, essential for embryonic development. Genes Dev. 10: 3141–3155.
- Amali, A.A., Rekha, R.D., Lin, C.J., Wang, W.L., Gong, H.Y., Her, G.M., and Wu, J.L. (2006). Thioacetamide induced liver damage in zebrafish embryo as a disease model for steatohepatitis. J. Biomed. Sci. 13: 225–232.
- Biemar, F., Argenton, F., Schmidtke, R., Epperlein, S., Peers, B., and Driever, W. (2001). Pancreas development in zebrafish: Early dispersed appearance of endocrine hormone expressing cells and their convergence to form the definitive islet. Dev. Biol. 230: 189–203.

- Blouin, A., Bolender, R.P., and Weibel, E.R. (1977). Distribution of organelles and membranes between hepatocytes and nonhepatocytes in the rat liver parenchyma. A stereological study. J. Cell Biol. 72: 441–455.
- Bort, R., Signore, M., Tremblay, K., Martinez Barbera, J.P., and Zaret, K.S. (2006). Hex homeobox gene controls the transition of the endoderm to a pseudostratified, cell emergent epithelium for liver bud development. Dev. Biol. 290: 44–56.
- Braunbeck, T., Gorge, G., Storch, V., and Nagel, R. (1990). Hepatic steatosis in zebra fish (Brachydanio rerio) induced by long-term exposure to gamma-hexachlorocyclohexane. Ecotoxicol. Environ. Saf. 19: 355–374.
- Chen, J., Ng, S.M., Chang, C.Q., Zhang, Z.H., Bourdon, J.C., Lane, D.P., and Peng, J.R. (2009). p53 isoform 113p53 is a p53 target gene that antagonizes p53 apoptotic activity via BdxL activation in zebrafish. Genes Dev. 23: 278–290.
- Chen, J., Ruan, H., Ng, S.M., Gao, C., Soo, H.M., Wu, W., Zhang, Z.H., Wen, Z.L., Lane, D.P., and Peng, J.R. (2005). Loss of function of def selectively up-regulates 113p53 expression to arrest expansion growth of digestive organs in zebrafish. Genes Dev. 19: 2900–2911.
- Cheng, W., Guo, L., Zhang, Z.H., Soo, H.M., Wen, C.M., Wu, W., and Peng, J.R. (2006). HNF factors form a network to regulate liver-enriched genes in zebrafish. Dev. Biol. 294: 482–496.
- Chocron, S., Verhoeven, M.C., Rentzsch, F., Hammerschmidt, M., and Bakkers, J. (2007). Zebrafish Bmp4 regulates left-right asymmetry at two distinct developmental time points. Dev. Biol. 305: 577–588.
- Chung, W.S., Shin, C.H., and Stainier, D.Y.R. (2008). Bmp2 signaling regulates the hepatic versus pancreatic fate decision. Dev. Cell 15: 738–748.
- Chung, W.S., and Stainier, D.Y.R. (2008). Intra-endodermal interactions are required for pancreatic beta cell induction. Dev. Cell 14: 582–593.
- Concordet, J.P., and Ingham, P. (1994). Developmental biology—catch of the decade. Nature 369: 19–20.
- Detrich, H.W., Kieran, M.W., Chan, F.Y., Barone, L.M., Yee, K., Rundstadler, J.A., Pratt, S., Ransom, D., and Zon, L.I. (1995). Intraembryonic hematopoietic cell migration during vertebrate development. Proc. Natl. Acad. Sci. USA 92: 10713–10717.
- Deutsch, G., Jung, J.N., Zheng, M.H., Lora, J., and Zaret, K.S. (2001). A bipotential precursor population for pancreas and liver within the embryonic endoderm. Development 128: 871–881.
- Dienstag, J.L. (2008). Hepatitis B virus infection. N Engl. J. Med. 359: 1486–1500.
- Dong, P.D.S., Munson, C.A., Norton, W., Crosnier, C., Pan, X.F., Gong, Z.Y., Neumann, C.J., and Stainier, D.Y.R. (2007). Fgf10 regulates hepatopancreatic ductal system patterning and differentiation. Nat. Genet. 39: 397–402.
- Douarin, N.M. (1975). An experimental analysis of liver development. Med. Biol. 53: 427–455.
- Driever, W., SolnicaKrezel, L., Schier, A.F., Neuhauss, S.C.F., Malicki, J., Stemple, D.L., Stainier, D.Y.R., Zwartkruis, F., Abdelilah, S., Rangini, Z., Belak, J., and Boggs, C. (1996) A genetic screen for mutations affecting embryogenesis in zebrafish. Development 123: 37–46.
- Duncan, S.A. (2003). Mechanisms controlling early development of the liver. Mech. Dev. 120: 19–33.
- Farooq, M., Sulochana, K.N., Pan, X.F., To, J.W., Sheng, D., Gong, Z.Y., and Ge, R.W. (2008). Histone deacetylase 3 (hdac3) is specifically required for liver development in zebrafish. Dev. Biol. 317: 336–353.
- Field, H.A., Ober, E.A., Roeser, T., and Stainier, D.Y.R. (2003). Formation of the digestive system in zebrafish. I. Liver morphogenesis.

Dev. Biol. 253: 279–290.

- Galarneau, L., Pare, J.F., Allard, D., Hamel, D., Levesque, L., Tugwood, J.D., Green, S., and Belanger, L. (1996). The alpha(1)-fetoprotein locus is activated by a nuclear receptor of the Drosophila FTZ-F1 family. Mol. Cell. Biol. 16: 3853–3865.
- Godinho, L., Mumm, J.S., Williams, P.R., Schroeter, E.H., Koerber, A., Park, S.W., Leach, S.D., and Wong, R.O.L. (2005). Targeting of amacrine cell neurites to appropriate synaptic laminae in the developing zebrafish retina. Development 132: 5069–5079.
- Grunwald, D.J., and Eisen, J.S. (2002). Headwaters of the zebrafishemergence of a new model vertebrate. Nat. Rev. Genet. 3: 717–724.
- Gualdi, R., Bossard, P., Zheng, M.H., Hamada, Y., Coleman, J.R., and Zaret, K.S. (1996). Hepatic specification of the gut endoderm in vitro: Cell signaling and transcriptional control. Genes Dev. 10: 1670–1682.
- Haffter, P., Granato, M., Brand, M., Mullins, M.C., Hammerschmidt, M., Kane, D.A., Odenthal, J., van Eeden, F.J.M., Jiang, Y.J., Heisenberg, C.P., Kelsh, R.N., Furutani-Seiki, M., Vogelsang, E., Beuchle, D., Schach, U., Fabian, C., and NussleinVolhard, C. (1996). The identification of genes with unique and essential functions in the development of the zebrafish, Danio rerio. Development 123: 1–36.
- Her, G.M., Chiang, C.C., Chen, W.Y., and Wu, J.L. (2003). In vivo studies of liver-type fatty acid binding protein (L-FABP) gene expression in liver of transgenic zebrafish (Danio rerio). FEBS Lett. 538: 125–133.
- Holtzinger, A., and Evans, T. (2005). Gata4 regulates the formation of multiple organs. Development 132: 4005–4014.
- Horne-Badovinac, S., Lin, D., Waldron, S., Schwarz, M., Mbamalu, G., Pawson, T., Jan, Y.N., Stainier, D.Y.R., and Abdelilah-Seyfried, S. (2001). Positional cloning of heart and soul reveals multiple roles for PKC lambda in zebrafish organogenesis. Curr. Biol. 11: 1492–1502.
- Horne-Badovinac, S., Rebagliati, M., and Stainier, D.Y.R. (2003). A cellular framework for gut-looping morphogenesis in zebrafish. Science 302: 662–665.
- Huang, H., Ruan, H., Aw, M.Y., Hussain, A., Guo, L., Gao, C., Qian, F., Leung, T., Song, H., Kimelman, D., Wen, Z., and Peng, J. (2008). Mypt1-mediated spatial positioning of Bmp2-producing cells is essential for liver organogenesis. Development 135: 3209–3218.
- Jin, H., Sood, R., Xu, J., Zhen, F., English, M.A., Liu, P.P., and Wen, Z.L. (2009). Definitive hematopoietic stem/progenitor cells manifest distinct differentiation output in the zebrafish VDA and PBI. Development 136: 647–654.
- Jung, J.N., Zheng, M.H., Goldfarb, M., and Zaret, K.S. (1999). Initiation of mammalian liver development from endoderm by fibroblast growth factors. Science 284: 1998–2003.
- Kahn, P. (1994). Zebrafish hit the big time. Science 264: 904–905.
- Kawamoto, S., Matsumoto, Y., Mizuno, K., Okubo, K., and Matsubara, K. (1996). Expression profiles of active genes in human and mouse livers. Gene 174: 151–158.
- Korzh, S., Emelyanov, A., and Korzh, V. (2001). Developmental analysis of ceruloplasmin gene and liver formation in zebrafish. Mech. Dev. 103: 137–139.
- Lam, S.H., Wu, Y.L., Vega, V.B., Miller, L.D., Spitsbergen, J., Tong, Y., Zhan, H.Q., Govindarajan, K.R., Lee, S., Mathavan, S., Murthy, K.R.K., Buhler, D.R., Liu, E.T., and Gong, Z.Y. (2006). Conservation of gene expression signatures between zebrafish and human liver tumors and tumor progression. Nat. Biotechnol. 24: 73–75.
- Lee, C.S., Friedman, J.R., Fulmer, J.T., and Kaestner, K.H. (2005a). The initiation of liver development is dependent on Foxa transcription factors. Nature 435: 944–947.
- Lee, Y., Grill, S., Sanchez, A., Murphy-Ryan, M., and Poss, K.D.

(2005b). Fgf signaling instructs position-dependent growth rate during zebrafish fin regeneration. Development 132: 5173–5183.

- Li, M., Xie, Y.H., Kong, Y.Y., Wu, X., Zhu, L., and Wang, Y. (1998). Cloning and characterization of a novel human hepatocyte transcription factor, hB1F, which finds and activates enhancer II of hepatitis B virus. J. Biol. Chem. 273: 29022–29031.
- Li, Z., Korzh, V., and Gong, Z.Y. (2007). Localized rbp4 expression in the yolk syncytial layer plays a role in yolk cell extension and early liver development. BMC Dev. Biol. 7: 117.
- Lin, W.W., Wang, H.W., Sum, C., Liu, D., Hew, C.L., and Chung, B.C. (2000). Zebrafish ftz-f1 gene has two promoters, is alternatively spliced, and is expressed in digestive organs. Biochem. J. 348: 439–446.
- Lorent, K., Yeo, S.Y., Oda, T., Chandrasekharappa, S., Chitnis, A., Matthews, R.R., and Pack, M. (2004). Inhibition of Jagged-mediated Notch signaling disrupts zebrafish biliary development and generates multi-organ defects compatible with an Alagille syndrome phenocopy. Development 131: 5753–5766.
- Matsumoto, K., Yoshitomi, H., Rossant, J., and Zaret, K.S. (2001). Liver organogenesis promoted by endothelial cells prior to vascular function. Science 294: 559–563.
- Matthews, R.P., Lorent, K., Manoral-Mobias, R., Huang, Y.H., Gong, W.L., Murray, I.V.J., Blair, I.A., and Pack, M. (2009). TNF alphadependent hepatic steatosis and liver degeneration caused by mutation of zebrafish S-adenosylhomocysteine hydrolase. Development 136: 865–875.
- Matthews, R.P., Lorent, K., Russo, P., and Pack, M. (2004). The zebrafish onecut gene hnf-6 functions in an evolutionarily conserved genetic pathway that regulates vertebrate biliary development. Dev. Biol. 274: 245–259.
- Mayer, A.N., and Fishman, M.C. (2003). nil per os encodes a conserved RNA recognition motif protein required for morphogenesis and cytodifferentiation of digestive organs in zebrafish. Development 130: 3917–3928.
- Mullins, M.C., Hammerschmidt, M., Haffter, P., and Nussleinvolhard, C. (1994). Large-scale mutagenesis in the zebrafish—in search of genes-controlling development in a vertebrate. Curr. Biol. 4: 189–202.
- Noel, E.S., Casal-Sueiro, A., Busch-Nentwich, E., Verkade, H., Dong, P.D.S., Stemple, D.L., and Ober, E.A. (2008). Organ-specific requirements for Hdac1 in liver and pancreas formation. Dev. Biol. 322: 237–250.
- Ober, E.A., Olofsson, B., Makinen, T., Jin, S.W., Shoji, W., Koh, G.Y., Alitalo, K., and Stainier, D.Y.R. (2004). Vegfc is required for vascular development and endoderm morphogenesis in zebrafish. EMBO Rep. 5: 78–84.
- Ober, E.A., Verkade, H., Field, H.A., and Stainier, D.Y.R. (2006). Mesodermal Wht2b signalling positively regulates liver specification. Nature 442: 688–691.
- Pack, M., SolnicaKrezel, L., Malicki, J., Neuhauss, S.C.F., Schier, A.F., Stemple, D.L., Driever, W., and Fishman, M.C. (1996). Mutations affecting development of zebrafish digestive organs. Development 123: 321–328.
- Passeri, M.J., Cinaroglu, A., Gao, C., and Sadler, K.C. (2009). Hepatic steatosis in response to acute alcohol exposure in zebrafish requires sterol regulatory element binding protein activation. Hepatology 49: 443–452.
- Pyati, U.J., Cooper, M.S., Davidson, A.J., Nechiporuk, A., and Kimelman, D. (2006). Sustained Bmp signaling is essential for cloaca development in zebrafish. Development 133: 2275–2284.
- Pyati, U.J., Webb, A.E., and Kimelman, D. (2005). Transgenic zebrafish reveal stage-specific roles for Bmp signaling in ventral and posterior mesoderm development. Development 132: 2333–2343.

- Rai, K., Chidester, S., Zavala, C.V., Manos, E.J., James, S.R., Karpf, A.R., Jones, D.A., and Cairns, B.R. (2007). Dnmt2 functions in the cytoplasm to promote liver, brain, and retina development in zebrafish. Genes Dev. 21: 261–266.
- Rai, K., Nadauld, L.D., Chidester, S., Manos, E.J., James, S.R., Karpf, A.R., Cairns, B.R., and Jones, D.A. (2006). Zebra fish Dnmt1 and Suv39h1 regulate organ-specific terminal differentiation during development. Mol. Cell. Biol. 26: 7077–7085.
- Reimold, A.M., Etkin, A., Clauss, I., Perkins, A., Friend, D.S., Zhang, J., Horton, H.F., Scott, A., Orkin, S.H., Byrne, M.C., Grusby, M.J., and Glimcher, L.H. (2000). An essential role in liver development for transcription factor XBP-1. Genes Dev. 14: 152–157.
- Reiter, J.F., Alexander, J., Rodaway, A., Yelon, D., Patient, R., Holder, N., and Stainier, D.Y.R. (1999). Gata5 is required for the development of the heart and endoderm in zebrafish. Genes Dev. 13: 2983–2995.
- Reiter, J.F., Kikuchi, Y., and Stainier, D.Y.R. (2001). Multiple roles for Gata5 in zebrafish endoderm formation. Development 128: 125–135.
- Rossi, J.M., Dunn, N.R., Hogan, B.L.M., and Zaret, K.S. (2001). Distinct mesodermal signals, including BMPs from the septum transversum mesenchyme, are required in combination for hepatogenesis from the endoderm. Genes Dev. 15: 1998–2009.
- Sadler, K.C., Amsterdam, A., Soroka, C., Boyer, J., and Hopkins, N. (2005). A genetic screen in zebrafish identifies the mutants vps18, nf2 and foie gras as models of liver disease. Development 132: 3561–3572.
- Sadler, K.C., Krahn, K.N., Gaur, N.A., and Ukomadu, C.