

# Molecular characteristics and virulence potential of *Listeria monocytogenes* isolates from Chinese food systems

Jianshun Chen<sup>a</sup>, Xiaokai Luo<sup>a</sup>, Lingli Jiang<sup>a</sup>, Peijie Jin<sup>a</sup>, Wei Wei<sup>a</sup>, Dongyou Liu<sup>b</sup>, Weihuan Fang<sup>a,\*</sup>

<sup>a</sup>Institute of Preventive Veterinary Medicine and Provincial Key Laboratory of Preventive Veterinary Medicine, Zhejiang University, Hangzhou, China

<sup>b</sup>College of Veterinary Medicine, Mississippi State University, MS, USA

## ARTICLE INFO

### Article history:

Received 30 April 2008

Received in revised form 12 August 2008

Accepted 16 August 2008

Available online 18 October 2008

### Keywords:

*Listeria monocytogenes*

Subtype

Internalin gene profiling

Virulence potential

Chinese food systems

## ABSTRACT

In this study, we examined *Listeria monocytogenes* isolates from Chinese food sources in an attempt to gain further insights on the molecular characteristics and virulence potential of this important foodborne pathogen. Of the 88 *L. monocytogenes* food isolates recovered, 42 (47.7%) were of serovars 1/2a or 3a; 23 (26.1%) of serovars 1/2b or 3b; 15 (17.0%) of 1/2c or 3c; 6 (6.8%) of serovars 4b, 4d or 4e; and 2 (2.2%) of serovars 4a or 4c. In contrast to *inlAB* locus conserved in all serovars, internalin cluster between *ascB* and *dapE* varies with different serovars, with *inlC2DE*, *inlGC2DE* and *inlGHE* predominantly in serovars 1/2b or 4b, serovar 1/2a and serovar 1/2c. While *inlF* existed in all the *inlGHE*- and *inlGC2DE*-containing isolates but 17.4% of those having *inlC2DE*, *lmo2026* existed in all the *inlGHE*-containing isolates but 20.0% of those bearing *inlGC2DE*, suggesting that *inlF* might have co-evolved with *inlGC2DE* and *inlGHE* while *lmo2026* with *inlGHE* only. With the exception of serovar 4a isolate, most serovar isolates demonstrated remarkable ability to form plaques on L929 cells and produced significant mouse mortality irrespective of the internalin gene organization and whether an intact *actA* gene is present or not. These results indicate that majority of these food isolates may have the potential to cause human diseases if ingested via contaminated foods. Given that serovar 4b accounts for nearly half of human clinical listeriosis cases documented, the relative low proportion of serovar 4b food isolates suggests that this serovar is probably more tolerant of the adverse conditions in the host's stomach and/or more efficient in entering host cells than serovars 1/2a, 1/2b and 1/2c.

© 2008 Elsevier Ltd. All rights reserved.

## 1. Introduction

*Listeria monocytogenes* is an intracellular bacterium that has the capability to infect a range of cell types, including professional phagocytes and non-phagocytes (e.g. epithelial cells, endothelial cells, hepatic cells and fibroblasts), and to cross the intestinal, blood-brain and placental barriers. Due to its widespread nature and its ability to tolerate wide pH, temperature and salt ranges, *L. monocytogenes* readily enters food processing facilities and survives and grows in a variety of food stuffs such as milk, seafood, vegetables and meat products. Accordingly, this bacterium has been responsible for an increasing proportion of human foodborne diseases, especially in infants, pregnant women, the elderly and immunosuppressed individuals, with a mortality rate approaching to 30% (Vazquez-Boland et al., 2001; Liu et al., 2006a,b).

*L. monocytogenes* infection process comprises several distinct stages: adhesion and invasion of host cells, escape from vacuole, intracellular multiplication and intercellular spread (Vazquez-Boland et al., 2001). *L. monocytogenes* adheres to and invades host cells both passively through phagocytosis and actively through actions of internalins, a complex family of leucine-rich repeat-containing (LRR) proteins (Bierne et al., 2007). Of the 25 members in the multigene internalin family, InlA binds to host adhesion protein E-cadherin for entering into epithelial cells and InlB interacts with a wide range of cell receptors for gaining entry into other cell types (Bergmann et al., 2002; Geese et al., 2002). InlC has been shown to be involved in the post-intestinal stage of infection (Engelbrecht et al., 1996), and InlJ plays a role in assisting *L. monocytogenes* crossing the intestinal barrier and also functions as an adhesin (Sabet et al., 2005, 2008). The genes encoding InlC and InlJ have recently been proved to be excellent markers for rapid determination of *L. monocytogenes* virulence (Liu et al., 2003, 2007a,b). In addition, *inlH*, which is contiguous with *inlG* and *inlE* between *ascB* and *dapE*, is also involved in listerial virulence (Dramsi et al., 1997; Bierne et al., 2007). Remarkably, some strains carry clusters *inlGC2DE* or *inlC2DE* in this region, with *inlC2* and

*inlD* instead of *inlH* (Nelson et al., 2004; Tsai et al., 2006). In contrast, *inlF* (which mediates invasion and virulence under specific conditions) and *lmo2026* (which possibly affects listerial multiplication in the brain) are located discretely in chromosomal regions (Bierne et al., 2007; Kirchner and Higgins, 2008). Once inside host cells, *L. monocytogenes* readily escapes from vacuole, replicates in the cytoplasm, and spreads to the neighboring cells to initiate a new infection cycle. Here, a *L. monocytogenes* surface protein ActA encoded by *actA* contributes significantly to the actin-based intra- and intercellular mobility in addition to being an adhesion factor (Suarez et al., 2001).

Because the usual route of *L. monocytogenes* infection in humans is via contaminated food, we conducted a survey of *L. monocytogenes* isolates recovered from Chinese food systems, and examined 12 internalin genes as well as *actA* gene in these isolates by PCR and DNA sequencing. This was followed by evaluation of selected *L. monocytogenes* food isolates for *in vitro* adhesion and cell-plaque formation as well as *in vivo* mouse virulence.

## 2. Materials and methods

### 2.1. Bacterial strains and DNA manipulations

A total of 88 *L. monocytogenes* isolates from various food sources in southeastern China were analyzed. These included 9 from milk products, 20 from seafood, 3 from vegetables, 22 from pork, 21 from chicken, 4 from beef, 3 from mutton, 3 from duck, 2 from cony meat and 1 from venison (Table 1). The food samples were homogenized and *Listeria* bacteria were enriched in 225 ml half Fraser broth for 4 h at 30 °C followed by full Fraser broth for 48 h at 37 °C before verification with *Listeria* selective agar plate (CHROMAgar *Listeria*, Paris, France) and multiplex PCR (Zeng et al., 2006). Additionally, 10 *L. monocytogenes* strains representing serovars 1/2a, 1/2b, 1/2c, 4a and 4b from reference collections were examined as controls in the study (Table 1). While the 88 *L. monocytogenes* food isolates were derived from different samples of given food types, there was a possibility that some isolates might belong to similar clones as they were all originated from a relatively small geographic region (surrounding Hangzhou city) in southeast China.

*Listeria* strains were retrieved from frozen glycerol stocks and cultured on brain heart infusion broth (BHI; Oxoid, Hampshire, England) at 37 °C. Genomic DNA was extracted from these strains using a protocol reported previously (Jiang et al., 2008). Briefly, *Listeria* cell pellet from 5 ml BHI culture was resuspended in 1 ml lysis buffer consisting of 2.0% Triton X-100 plus 2.5 mg sodium azide/ml in 0.1 M Tris-HCl buffer at pH 8.0 and proteinase K (10 µl of 20 mg/ml stock). After incubation at 55 °C overnight, *Listeria* DNA was extracted with phenol-chloroform and precipitated in isopropanol. Oligonucleotide primers were synthesized by Invitrogen Biotechnology (Shanghai, China) (Table 2), and Taq DNA polymerase (TaKaRa Biotech Co. Ltd., Dalian, China) was used for PCR amplification. PCR was conducted using a thermal cycler (MJ Research Inc. MA, Boston, U.S.A.), with annealing temperatures depending on specific primer pairs (Table 2), and the duration of extension depending on the expected length of amplicon (1 min per kb, at 72 °C). For DNA sequencing analysis, PCR fragments were purified with the AxyPrep DNA Gel Extraction Kit (Axygen Inc., USA) and their sequences determined by dideoxy method on ABI-PRISM 377 DNA sequencer.

### 2.2. Phylogenetic analysis of *actA* gene

A 537 bp (or 432 bp in some cases) fragment (corresponding to the *actA* nucleotide positions between 775 and 1313, which covers four proline-rich repeats or PRRs) was amplified from the 88 *L. monocytogenes* food isolates as well as reference strains by PCR

using *actA* gene primers and its nucleotide sequence analyzed (Table 2). The deduced amino acid sequences were aligned for lineage classification (Wiedmann et al., 1997) by using the Molecular Evolutionary Genetics Analysis software (MEGA version 3.0) (<http://www.megasoftware.net>).

### 2.3. Serotype identification

A multiplex PCR with primers from ORF2819, ORF2110, *lmo0737* and *lmo1118* was performed for typing the food isolates (Doumith et al., 2004a). Specifically, ORF2819 primers recognize serovars 1/2b, 3b, 4b, 4d and 4e; ORF2110 primers further separate serovars 4b, 4d and 4e from serovars 1/2b and 3b; *lmo0737* primers identify serovars 1/2a, 3a, 1/2c and 3c strains; and *lmo1118* further distinguish serovars 1/2c and 3c from 1/2a and 3a (Doumith et al., 2004a). Also, *lmo1134* primers with specificity for all *L. monocytogenes* strains except serovars 4a and 4c were utilized (Liu et al., 2003). Serotyping based on agglutination between somatic/flagellar antigens and specific antisera (Schonberg et al., 1996) was applied to selected *L. monocytogenes* isolates, including all from milk and seafood, and several from vegetable, pork and chicken along with the reference strains, to confirm the accuracy of serogrouping results based on multiplex PCR.

### 2.4. Examination of internalin genes

As the internalin genes are known to be scattered in different sections of *L. monocytogenes* genome and contribute to bacterial adhesion and/or virulence (Bierne et al., 2007), we investigated the presence or absence of 12 internalin genes in the 88 *L. monocytogenes* food isolates by PCR with specific primers outlined in Table 2. Due to the conserved repeats present in internalin multigene family (Bierne et al., 2007), primers were designed based on the distinguishable regions through sequence comparison. As *inlH* and *inlC2* shared highly identical nucleotide sequences, a common primer set was employed. To further examine the internalin organization between *ascB* and *dapE*, upstream primer (u, targeting *inlG*) was combined with downstream primer (d1, targeting *inlE*; d2, targeting *inlD*) respectively (Table 2). Bridging PCR using primer pair u/d1 was expected to produce 4000 bp fragment from strains harboring *inlGC2DE*, 2241 bp fragment from strains harboring *inlGHE* cluster and no fragment from those harboring *inlC2DE* or being empty between *ascB* and *dapE*, while PCR using primer pair u/d2 only yield 2241 bp fragment from strains harboring *inlGC2DE*.

### 2.5. Adhesion assay

The ability of selected *L. monocytogenes* isolates to adhere to HeLa epithelial cells was examined (Olier et al., 2002; Jiang et al., 2006). As for plaque-forming and mouse virulence assays, food isolates were selected for adhesion assay on the basis of their distinct or unusual internalin and *actA* gene profiles in addition to the control strains. Briefly, HeLa monolayers at 80% confluence in Dulbecco's modified Eagle's medium (DMEM) (Gibco, USA) in 12-well plates (Corning, USA) were inoculated with 300 µl of fresh bacterial suspension ( $10^7$  colony forming units or CFU per ml) to obtain a multiplicity of infection (MOI) of 10:1 for 1 h at 37 °C in the presence of 5% CO<sub>2</sub>. The cell monolayers were then washed three times with 10 mM phosphate-buffered saline (PBS) pH 7.2 to remove nonadherent bacteria. Adherent bacteria were harvested after lysis of the cell monolayers with 250 µl Triton X-100 (0.25% in cold PBS) and 250 µl trypsin for 10 min at 4 °C. The CFU values for viable bacteria were determined by plating suitable dilutions of the lysates onto BHI plates. The plates were subsequently incubated for 36 h at 37 °C. We set the adhesion rate of the serovar 4b strain M5

**Table 1**  
Characteristics of *L. monocytogenes* reference and food isolates under investigation.

Stain/isolate	Source	Lineage <sup>a</sup>	Multiplex PCR <sup>b</sup>	Agglutination <sup>c</sup>	actA <sup>a</sup>	inA	inB	inC	inD	inE	inF	inG	inH/C2	inI	inJ	lmo2026	inG-E <sup>d</sup> (bp)	inG-D <sup>e</sup> (bp)	ascB dapE structure
ScottA	Reference	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
NICPBP 54007	Reference	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
SLCC2755	Reference	I	1/2b or 3b	1/2b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
EGD-e	Reference	II	1/2a or 3a	1/2a	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
10403S	Reference	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
NICPBP 54003	Reference	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
AB2483	Reference	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
NICPBP 54002	Reference	II	1/2c or 3c	1/2c	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
NCTC 5348	Reference	II	1/2c or 3c	1/2c	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
NICPBP 54006	Reference	III	4a or 4c	4a	432	+	+	-	-	-	-	-	-	-	-	-	-	-	-
M1	Milk	I	1/2b or 3b	1/2b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
M2	Milk	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
M3	Milk	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
M4	Milk	II	1/2a, or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
M5	Milk	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
M6	Milk	II	1/2a, or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
M7	Milk	III	4a or 4c	4a	432	+	+	-	-	-	-	-	-	-	-	-	-	-	-
M8	Milk	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	-	+	+	+	-	-	-	inIC2DE
M9	Milk	II	1/2c or 3c	1/2c	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
S1	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
S2	Seafood	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S3	Seafood	II	1/2c or 3c	1/2c	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
S4	Seafood	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S5	Seafood	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S6	Seafood	I	4b,4d or 4e	4ab	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S7	Seafood	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S8	Seafood	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S9	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
S10	Seafood	I	1/2b or 3b	1/2b	432	-	-	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S11	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S12	Seafood	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S13	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
S14	Seafood	I	1/2b or 3b	1/2b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
S15	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
S16	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
S17	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
S18	Seafood	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
S19	Seafood	III	4a or 4c	4b	537	+	+	+	+	+	-	-	+	+	+	-	4000	2241	inIGC2DE
S20	Seafood	II	1/2c or 3c	1/2c	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
V1	Vegetable	II	1/2c or 3c	1/2c	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
V2	Vegetable	I	4b,4d or 4e	4b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
V3	Vegetable	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
P1	Pork	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
P2	Pork	I	1/2b or 3b	1/2b	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P3	Pork	II	1/2a or 3a	1/2a	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P4	Pork	I	4b,4d or 4e	4b	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P5	Pork	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P6	Pork	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	-	-	-	inIC2DE
P7	Pork	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P8	Pork	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
P9	Pork	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
P10	Pork	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P11	Pork	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
P12	Pork	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
P13	Pork	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P14	Pork	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
P15	Pork	I	1/2b or 3b	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P16	Pork	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P17	Pork	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
P18	Pork	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
P19	Pork	II	1/2c or 3c	1/2c	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
P20	Pork	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
P21	Pork	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
P22	Pork	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
C1	Chicken	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
C2	Chicken	II	1/2a or 3a	1/2a	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
C3	Chicken	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	inIGC2DE
C4	Chicken	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE
C5	Chicken	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
C6	Chicken	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
C7	Chicken	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	inIGC2DE
C8	Chicken	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
C9	Chicken	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	inIC2DE
C10	Chicken	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	inIGHE

(continued on next page)

Table 1 (continued)

Stain/isolate	Source	Lineage <sup>a</sup>	Multiplex PCR <sup>b</sup>	Agglutination <sup>c</sup>	<i>actA</i> <sup>a</sup>	<i>inA</i>	<i>inB</i>	<i>inC</i>	<i>inD</i>	<i>inE</i>	<i>inF</i>	<i>inG</i>	<i>inH/C2</i>	<i>inI</i>	<i>inJ</i>	<i>lmo2026</i>	<i>inG-E</i> <sup>d</sup> (bp)	<i>inG-D</i> <sup>e</sup> (bp)	<i>ascB</i> <i>dapE</i> structure
C11	Chicken	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	<i>inlGC2DE</i>
C12	Chicken	I	1/2b or 3b	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
C13	Chicken	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
C14	Chicken	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
C15	Chicken	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	<i>inlGHE</i>
C16	Chicken	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	<i>inlGHE</i>
C17	Chicken	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
C18	Chicken	II	1/2a or 3a	1/2a	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	<i>inlGC2DE</i>
C19	Chicken	II	1/2a or 3a	ND	537	+	+	+	+	+	+	-	+	+	+	-	-	-	<i>inlC2DE</i>
C20	Chicken	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
C21	Chicken	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
B1	Beef	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	<i>inlGC2DE</i>
B2	Beef	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
B3	Beef	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	+	4000	2241	<i>inlGC2DE</i>
B4	Beef	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
G1	Mutton	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
G2	Mutton	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	<i>inlGC2DE</i>
G3	Mutton	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
D1	Duck	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	<i>inlGC2DE</i>
D2	Duck	II	1/2c or 3c	ND	537	+	+	+	-	+	+	+	+	+	+	+	2241	-	<i>inlGHE</i>
D3	Duck	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
R1	Cony meat	I	1/2b or 3b	ND	537	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>
R2	Cony meat	II	1/2a or 3a	ND	537	+	+	+	+	+	+	+	+	+	+	-	4000	2241	<i>inlGC2DE</i>
E1	Vension	II	1/2a or 3a	ND	432	+	+	+	+	+	-	-	+	+	+	-	-	-	<i>inlC2DE</i>

<sup>a</sup> The lineage of *L. monocytogenes* strains/isolates was ascertained on the basis of the *actA* gene sequences.

<sup>b</sup> The multiplex PCR employs primers from ORF2819, ORF2110, *lmo0737* and *lmo1118*, and divides into five serotyping groups: 1/2a or 3a; 1/2b or 3b; 1/2c or 3c; 4b, 4d or 4e; 4a or 4c.

<sup>c</sup> The agglutination assay was performed as previously described by Schonberg et al. (1996).

<sup>d</sup> PCR employing primers u and d1 (Table 2) yield 4000 bp fragment from strains harboring *inlGC2DE*, 2241 bp fragment from strains harboring *inlGHE* or none from strains harboring *inlC2DE* or being empty between *ascB* and *dapE*.

<sup>e</sup> PCR employing primers u and d2 (Table 2) yield 2241 bp fragment from strains harboring *inlGC2DE* or none from other strains.

Table 2

Identity and sequence of PCR primers used in this study.

Gene	Specificity	Forward and reverse primers (5'-3')	Product size (bp)	Annealing temperature (°C)	Reference
<i>lmo0737</i>	<i>L. monocytogenes</i> serovars 1/2a, 3a, 1/2c and 3c	AGGGCTTCAAGGACTTACCC ACGATTTCTGCTTGCCATTC	691	53	Doumith et al., 2004a
<i>lmo1118</i>	<i>L. monocytogenes</i> serovars 1/2c and 3c	AGGGGTCTTAAATCTGGAA CGGCTTGTTCGGCATACTTA	906	53	Doumith et al., 2004a
ORF2819	<i>L. monocytogenes</i> serovars 1/2b, 3b, 4b, 4d, 4e and 7	AGCAAATGCCAAAATCTCGT CATCACTAAAGCCTCCCATTC	471	53	Doumith et al., 2004a
ORF2110	<i>L. monocytogenes</i> serovars 4b, 4d and 4e	AGTGGACAATTGATTGGTGA CATCCATCCCTTACTTTGGAC	597	53	Doumith et al., 2004a
<i>lmo1134</i>	All <i>L. monocytogenes</i> serovars but 4a, 4c and 7	ACCCGATAGCAAGGAGGAAC AACTTCTCTCGATACCCATCCA	367	53	Liu et al., 2003
<i>actA</i>	All <i>L. monocytogenes</i> serovars	GGTACGTGATAAAATCGACGA TAGTTATGTCACITTCAGAGC	537 or 432 <sup>a</sup>	55	Wiedmann et al., 1997
<i>inA</i>	All <i>L. monocytogenes</i> serovars	TAATATAAGTGATATAAGCCCAG TTTATCCGTACTGAAATTC	606	60	This study
<i>inB</i>	All <i>L. monocytogenes</i> serovars	ATCACTTTCTTTGGAGCATAATGGT GCCATCATCACTTATTATTCTGGA	394	60	This study
<i>inC</i>	All <i>L. monocytogenes</i> but serovar 4a and some 4c	CCATCTGGTCTTTGACAGTA CAAATAAGTGACCTTAGTCCTT	398	55	This study
<i>inD</i>	Some lineages I and II strains	CTGTAGTAATGGCAATTAGCCT TGTTATTAGGACCACAAGCT	870	52	This study
<i>inE</i>	All <i>L. monocytogenes</i> but serovar 4a and some 4c	AGCTCAAAAGAAGTACAAGCA GTGCAATAAGCTCACCAGAAA	787	55	This study
<i>inF</i>	Some lineage II strains	TGACTTATTTCAGTTGGGGT TTGGTTCAGGAATAAGCGCG	1119	55	This study
<i>inG</i>	Some lineage II strains	GTGAAGACGGAACCTTGAAA GCTTCTACTATCGGTTGAACA	668	52	This study
<i>inH/C2</i>	All <i>L. monocytogenes</i> but serovar 4a and some 4c	ATAGCTACTTTATCAGCATTT ATATCACTTATTTATTATCATC	437	52	This study
<i>inI</i>	All <i>L. monocytogenes</i> but serovar 4a and some 4c	GTTTCCAGACGACAATCTTGCTA AATCGGTACAGTTACTCGCATCA	635	58	This study
<i>inJ</i>	All <i>L. monocytogenes</i> but serovar 4a and some 7	TAGATGTGACACCACAACTCAA TGTATTATCGGTGACATCAAGCT	401	58	This study
<i>lmo2026</i>	Some lineage II strains	CGGTTGTTCCTGATGTGTTGCTT TCGACGAACCTAATCCTTTTGC	837	58	This study
<i>ascB</i> <i>dapE</i> region	All <i>L. monocytogenes</i> but serovar 4a and some 4c	u-TGATGATTAAGTATGATTCTTA d1-ATCAGTAAGCACTGGATCAGTA d2-CGTTTGCTAAATTCATCTGTGA	Variable <sup>b</sup>	55	This study

<sup>a</sup> Some *L. monocytogenes* strains exhibit a deletion of 105 nucleotides in *actA* gene, leading to removal of 35 amino acids in the ActA protein.

<sup>b</sup> Primer pairs u/d1 and u/d2 both yield variable product sizes from different strains (for more information see Table 1).

at 100%. For statistical analysis, the two-tailed Student's *t*-test was applied and *P*-values of  $\leq 0.05$  were considered as statistically significant.

## 2.6. Plaque-forming assay

The ability of selected *L. monocytogenes* isolates to form plaques on mouse fibroblasts L929 cells was assessed (Roberts et al., 2005; Jiang et al., 2006). Cell monolayers were grown to 80% confluence in 2 ml DMEM containing 10% fetal bovine serum in six-well plates (Corning, USA). The overnight *Listeria* cultures were centrifuged and resuspended in PBS. For each strain tested, one well was infected with  $5 \times 10^5$  CFU and the other was infected with  $1.5 \times 10^5$  CFU. Upon 1-h incubation at 37 °C, the cell monolayers were washed three times with PBS and overlaid with 3 ml of DMEM containing 20 µg/ml gentamicin and 1.4% agarose (Oxoid Ltd., Hampshire, England). Following 3-day incubation at 37 °C, a second 2-ml overlay of DMEM containing 0.02% neutral red solution and 1.4% agarose was added. After a final day of incubation, plaques were photographed by a digital camera. The diameters of 25 plaques were measured using adobe photoshop software for each strain. The plaque size of serovar 4b strain M5 was set at 100%.

## 2.7. Virulence in mice

The ability of selected *L. monocytogenes* isolates to cause pathogenic effects on mice was assessed (Jiang et al., 2006, 2007). Five groups (six per group) of female ICR mice at 20–22 g (Zhejiang College of Traditional Chinese Medicine, Hangzhou, China) were inoculated intraperitoneally with 0.2 ml aliquots of appropriately diluted *Listeria* strain resuspended in PBS. Mice in the control group were injected with 0.2 ml PBS. The LD<sub>50</sub> values for mice were calculated by using the trimmed Spearman–Karber method on the basis of mouse mortality data recorded during a ten-day post-injection period. Relative virulence (%) of *L. monocytogenes* isolates

L.ser0s0,List-303T101Tf18.8(gr.819667TD(2-ml).801(ViruCr)-ohday)ydroshoe07 3erThe aeriity o0(oTJ/T111Tf18.8-184d(L.)-326(m54ocytog)10ts)-g



**Table 6**  
Relationship between *L. monocytogenes* *actA*, internalin gene status and *in vitro* and *in vivo* virulence.

Strain	Serovar	<i>actA</i> PCR (bp)	<i>inlC</i> PCR	<i>inlJ</i> PCR	<i>ascB</i> <i>dapE</i> structure	Relative adhesion rate $\pm$ SD <sup>a</sup> (%)	Relative size of plaque $\pm$ SD <sup>a</sup> (%)	Mice	
								log LD <sub>50</sub>	Relative virulence (%)
ScottA	4b	537	+	+	<i>inlC2DE</i>	ND	ND	5.50	60
NICPBP 54007	4b	537	+	+	<i>inlC2DE</i>	ND	ND	6.79	37
EGD-e	1/2a	537	+	+	<i>inlGHE</i>	ND	ND	6.64	37
10403S	1/2a	537	+	+	<i>inlGC2DE</i>	ND	92.3 $\pm$ 5.8	5.49	60
NICPBP 54006	4a	432	–	–	–	16.2 $\pm$ 2.2	ND	8.35	3
S5	1/2b	432	+	+	<i>inlC2DE</i>	ND	94.5 $\pm$ 3.5	5.94	53
S7	1/2b	432	+	+	<i>inlC2DE</i>	ND	84.9 $\pm$ 3.1	5.79	57
S8	1/2b	432	+	+	<i>inlC2DE</i>	47.6 $\pm$ 1.5	92.7 $\pm$ 2.2	5.08	70
M1	1/2b	537	+	+	<i>inlC2DE</i>	54.1 $\pm$ 4.8	94.9 $\pm$ 8.2	6.46	40
P2	1/2b	537	+	+	<i>inlC2DE</i>	ND	82.0 $\pm$ 1.3	6.45	43
C17	1/2b or 3b	537	+	+	<i>inlC2DE</i>	ND	ND	5.83	53
M2	1/2b	432	+	+	<i>inlC2DE</i>	ND	77.3 $\pm$ 0.4	6.43	40
M3	1/2b	432	+	+	<i>inlC2DE</i>	ND	89.6 $\pm$ 0.7	6.32	43
S10	1/2b	432	+	+	<i>inlC2DE</i>	6.7 $\pm$ 1.1	ND	ND	ND
S2	4b	537	+	+	<i>inlC2DE</i>	101.3 $\pm$ 5.2	78.0 $\pm$ 3.9	6.74	37
S4	4b	537	+	+	<i>inlC2DE</i>	ND	79.0 $\pm$ 4.5	6.72	37
M5	4b	537	+	+	<i>inlC2DE</i>	100 $\pm$ 0	100 $\pm$ 0	3.86	93
S6	4ab	537	+	+	<i>inlC2DE</i>	99.7 $\pm$ 10.5	83.2 $\pm$ 0.7	4.40	83
S9	1/2a	537	+	+	<i>inlGHE</i>	50.8 $\pm$ 3.2	91.2 $\pm$ 1.4	6.31	43
M4	1/2a	537	+	+	<i>inlGC2DE</i>	31.3 $\pm$ 3.2	82.5 $\pm$ 2.3	5.45	60
M6	1/2a	537	+	+	<i>inlGC2DE</i>	ND	88.4 $\pm$ 13.1	5.55	60
S1	1/2a	537	+	+	<i>inlGC2DE</i>	26.2 $\pm$ 3.7	95.8 $\pm$ 7.8	5.53	60
S17	1/2a	537	+	–	<i>inlGC2DE</i>	13.7 $\pm$ 6.0	ND	ND	ND
S18	1/2a	537	+	–	<i>inlGC2DE</i>	15.9 $\pm$ 3.8	ND	ND	ND
P1	1/2a	537	+	+	<i>inlGC2DE</i>	ND	88.9 $\pm$ 1.2	6.26	47
P3	1/2a	537	+	+	<i>inlC2DE</i>	23.7 $\pm$ 2.2	90.8 $\pm$ 3.4	6.07	50
C18	1/2a	537	+	+	<i>inlGC2DE</i>	34.4 $\pm$ 4.5	ND	ND	ND
S11	1/2a	537	+	+	<i>inlC2DE</i>	40.4 $\pm$ 3.0	ND	ND	ND
P6	1/2a	537	+	+	<i>inlC2DE</i>	46.6 $\pm$ 7.2	ND	ND	ND
C2	1/2a	432	+	+	<i>inlC2DE</i>	38.7 $\pm$ 4.8	ND	ND	ND
P19	1/2c	537	+	+	<i>inlGHE</i>	28.5 $\pm$ 3.1	ND	ND	ND
V1	1/2c	537	+	+	<i>inlGHE</i>	45.6 $\pm$ 3.6	78.5 $\pm$ 1.3	6.11	50
S3	1/2c	537	+	+	<i>inlGC2DE</i>	26.2 $\pm$ 3.2	84.9 $\pm$ 0.6	6.19	47
M7	4a	432	–	–	–	25.0 $\pm$ 3.1	0	8.21	7
S19	(4c)	537	+	+	<i>inlGC2DE</i>	65.6 $\pm$ 7.3	ND	6.83	33

ND, not done.

<sup>a</sup> The adhesion rate and plaque size of serovar 4b strain M5 was set at 100%.

and *inlJ*). On the other hand, the fact that five serovar 1/2b strains (S5, S7, S8, M2 and M3) with a 105-bp deletion in the *actA* gene had no difficulty spreading to other cells, and causing significant mouse mortality (with relative virulence of 40–70%) undermines the role of *actA* in listerial virulence and highlights the possible existence of other mechanisms for *L. monocytogenes* spreading than the requirement of a fully functional ActA protein. Indeed, a serovar 4b strain F2365 from Jalisco cheese outbreak of 1985 in California (Nelson et al., 2004) has been shown to contain a 105-bp deletion in its *actA* gene, which did not seem to stop the bacterium causing listeriosis in humans. As a number of internalins (e.g. *inlC*, *inlJ* and *lmo2470*), transcriptional regulators (e.g. *lmo1134*, *lmo1116* and *lmo2470*) are absent in serovar 4a strain M7, but present in other serovars (Liu et al., 2003, 2007a,b), the distinct genomic background including the presence of additional internalin and transcriptional regulator genes in *L. monocytogenes* serovar 1/2b strains S5, S7, S8, M2 and M3 may have contributed partially to their cell-to-cell spread and virulence.

Furthermore, the *inlAB* locus encodes critical proteins for *L. monocytogenes* entry into epithelial and other cells types (Bierne et al., 2007), and there is evidence that nucleotide alterations in this locus may reduce its virulence in mammalian hosts (Olier et al., 2002). Indeed, several *L. monocytogenes* strains producing truncated InlA protein have been isolated from human carriers, and these strains displayed reduced ability to cause disease in murine models (Olier et al., 2002). Isolation of a serovar 1/2b strain (S10) from seafood in this study represents the first instance of *L. monocytogenes* bacterium which loses the whole *inlAB* locus in the food environment.

Although the organization of internalin genes between *ascB* and *dapE* failed to further delineate listerial adhesion and intercellular abilities and virulence (Table 6), *ascB* *dapE* structure offered a potential marker for lineages and serovars from phylogenetic perspective. Clusters *inlC2DE*, *inlGC2DE* and *inlGHE* were supposed to be the ancestral versions of lineage I (including serovars 1/2b, 3b, 4b, 4d, and 4e), serovar 1/2a (or 3a) and serovar 1/2c (or 3c) respectively. Accordingly, serovar 1/2a (or 3a) strains carrying *inlGHE* (e.g. EGD-e and S9) and serovar 1/2c (or 3c) strains carrying *inlGC2DE* (e.g. NICPBP 54002, NCTC 5348 and S3) appeared as the atypical replacements. In fact, EGD-e was hinted to be an atypical serovar 1/2a strain in previous reports (Hain et al., 2007). Moreover, *inlF* existed in all the *inlGHE*-containing and *inlGC2DE*-containing isolates but 17.4% of *inlC2DE*-containing ones of lineage II, while *lmo2026* existed in all the *inlGHE*-containing isolates, 20.0% of *inlGC2DE*-containing ones but none of *inlC2DE*-containing isolates, suggesting that *inlF* might have co-evolved with *inlGC2DE* and *inlGHE* while *lmo2026* with *inlGHE* only.

If we consider gene deletion as an important force in *Listeria* evolution, *L. monocytogenes* serovar 1/2c (or 3c) strains, containing a complete set of internalins together with intact ActA (Table 3), is possibly more ancestral than those of other serovars, and evolved to serovars 1/2a (or 3a), 1/2b (or 3b), and 4b (or 4d, 4e and 4ab), and then to serovar 4a via gene deletion and/or horizontal gene transfer. Previously, upon examination of 264 genes from 113 *L. monocytogenes* strains by microarray, Doumith et al. (2004b) have also shown the possibility of *L. monocytogenes* serovars evolving from serovar 4b, to serovar 4c, and then to serovar 4a. Given the absence

of *inlGC2DE* and other genes in *Listeria innocua* in comparison with *L. monocytogenes* serovar 4a, the authors further hypothesized the transition from *L. monocytogenes* serovar 4a to *L. innocua*. At this stage, the whole genome sequences of *L. monocytogenes* serovars 1/2a and 4b, *L. innocua* and *Listeria welshimeri* (Glaser et al., 2001; Nelson et al., 2004; Hain et al., 2007) have been published, and the sequencing analyses of other *L. monocytogenes* serovars and *Listeria* species are in progress. There is no doubt that the evolutionary and phylogenetic links among *L. monocytogenes* serovars and *Listeria* species will be much clearer when the full genomic sequence data of representative strains become available in the near future.

In addition, while conventional agglutination assay is useful for serotyping *L. monocytogenes* strains, previous reports have indicated that this typing procedure is unable to correctly identify serovars 4a, 4b and 4c (Liu et al., 2006a,b; Schonberg et al., 1996). The fact that strain S19 was recognized as serovar 4b, but as 4c by other molecular typing techniques (e.g. *actA* gene sequence analysis and PCR targeting ORF2819, *lmo1134* and *inlJ* genes), reinforces again the continuing need to develop improved serotyping procedures for diagnostic and epidemiological applications. Toward this end, several lineage- and group-specific genes have been utilized for more precise serotyping of *L. monocytogenes* (Doumith et al., 2004a). Additionally, a multiplex PCR assay targeting *inlA*, *inlC* and *inlJ* genes has been devised for species- and virulence-specific determination (Liu et al., 2007b). The results from this study offer support for the use of *inlC* and *inlJ* as virulence markers since food strains harboring these genes were capable of causing severe mouse mortality via intraperitoneal route, whereas food strain M7 without these genes produced negligible mouse mortality.

## References

- Barbour, A.H., Rampling, A., Hormache, C.E., 2001. Variation in the infectivity of *Listeria monocytogenes* isolates following intragastric inoculation of mice. *Infect. Immun.* 69, 4657–4660.
- Bergmann, B., Raffelsbauer, D., Kuhn, M., Goetz, M., Hom, S., Goebel, W., 2002. InlA but not InlB-mediated internalization of *Listeria monocytogenes* by non-phagocytic mammalian cells needs the support of other internalins. *Mol. Microbiol.* 43, 557–570.
- Bierne, H., Sabet, C., Personnic, N., Cossart, P., 2007. Internalins: a complex family of leucine-rich repeat-containing proteins in *Listeria monocytogenes*. *Microbes Infect.* 9, 1156–1166.
- Chakraborty, T., Ebel, F., Wehland, J., Dufrenne, J., Notermans, S., 1994. Naturally occurring virulence-attenuated isolates of *Listeria monocytogenes* capable of inducing long term protection against infection by virulent strains of homologous and heterologous serotypes. *FEMS Immunol. Med. Microbiol.* 10, 1–9.
- Doumith, M., Buchrieser, C., Glaser, P., Jacquet, C., Martin, P., 2004a. Differentiation of the major *Listeria monocytogenes* serovars by multiplex PCR. *J. Clin. Microbiol.* 42, 3819–3822.
- Doumith, M., Cazalet, C., Simoes, N., Frangeul, L., Jacquet, C., Kunst, F., Martin, P., Cossart, P., Glaser, P., Buchrieser, C., 2004b. New aspects regarding evolution and virulence of *Listeria monocytogenes* revealed by comparative genomics and DNA arrays. *Infect. Immun.* 72, 1072–1083.
- Dramsi, S., Dehoux, P., Lebrun, M., Goossens, P.L., Cossart, P., 1997. Identification of four new members of the internalin multigene family of *Listeria monocytogenes* EGD. *Infect. Immun.* 65, 1615–1625.
- Engelbrecht, F., Chun, S.K., Osch, C., Hess, J., Lottspeich, F., Goebel, W., Sokolovic, A., 1996. A new PrfA-regulated gene of *Listeria monocytogenes* encoding a small, secreted protein which belongs to the family of internalins. *Mol. Microbiol.* 21, 823–837.
- Ermolaeva, S., Karpova, T., Novella, S., Wagner, M., Scotti, M., Tartakovskii, I., Vazquez-Boland, J.A., 2003. A simple method for the differentiation of *Listeria monocytogenes* on induction of lecithinase activity by charcoal. *Int. J. Food Microbiol.* 82, 87–94.
- Geese, M., Loureiro, J.J., Bear, J.E., Wehland, J., Gertler, F.B., Sechi, A.S., 2002. Contribution of Ena/VASP proteins to intracellular motility of *Listeria monocytogenes* phosphorylation and proline-rich core but not F-actin binding or multimerization. *Mol. Biol. Cell* 13, 2383–2396.
- Geoffroy, C., Gaillard, J.L., Alouf, J.E., Berche, P., 1989. Production of thiol-dependent haemolysins by *Listeria monocytogenes* and related species. *J. Gen. Microbiol.* 135, 481–487.
- Glaser, P., Frangeul, L., Buchrieser, C., Rusniok, C., Amend, A., Baquero, F., Berche, P., Bloecker, H., Brandt, P., Chakraborty, T., Charbit, A., Chetouani, F., Couve, E., de Daruvar, A., Dehoux, P., Domann, E., Dominguez-Bernal, G., Duchaud, E., Durant, L., Dussurget, O., Entian, K.D., Fsihi, H., Portillo, F.G., Garrido, P., Gautier, L., Goebel, W., Gomez-Lopez, N., Hain, T., Hauf, J., Jackson, D., Jones, L.M., Kaerst, U., Kreft, J., Kuhn, M., Kunst, F., Kurapatk, G., Madueno, E., Maitournam, A., Vicente, J.M., Ng, E., Nedjari, H., Nordsiek, G., Novella, S., de Pablos, B., Perez-Diaz, J.C., Purcell, R., Remmel, B., Rose, M., Schluter, T., Simoes, N., Tierrez, A., Vazquez-Boland, J.A., Voss, H., Wehland, J., Cossart, P., 2001. Comparative genomics of *Listeria* species. *Science* 294, 849–852.
- Goulet, V., Jacquet, C., Martin, P., Vaillant, V., Laurent, E., de Valk, H., 2006. Surveillance of human listeriosis in France, 2001–2003. *Euro Surveill.* 11, 79–81.
- Hain, T., Chatterjee, S.S., Ghai, R., Kuenne, C.T., Billionm, A., Steinweg, C., Domann, E., Karst, U., Jansch, L., Wehland, J., Eisenreich, W., Bacher, A., Joseph, B., Schar, J., Kreft, J., Klumpp, J., Loessner, M.J., Dorscht, J., Neuhaus, K., Fuchs, T.M., Scherer, S., Doumith, M., Jacquet, C., Martin, P., Cossart, P., Rusniok, C., Glaser, P., Buchrieser, C., Goebel, W., Chakraborty, T., 2007. Pathogenomics of *Listeria* spp. *Int. J. Med. Microbiol.* 297, 541–547.
- Hong, E., Doumith, M., Duperrier, S., Giovannacci, I., Morvan, A., Glaser, P., Buchrieser, C., Jacquet, C., Martin, P., 2007. Genetic diversity of *Listeria monocytogenes* recovered from infected persons and pork, seafood and dairy products on retail sale in France during 2000 and 2001. *Int. J. Food Microbiol.* 114, 187–194.
- Jaradat, Z., Bhunia, A.K., 2003. Adhesion, invasion and translocation characteristics of *Listeria monocytogenes* serotypes in Caco-2 cell and mouse models. *Appl. Environ. Microbiol.* 69, 3640–3645.
- Jiang, L., Xu, J., Chen, N., Shuai, J., Fang, W., 2006. Virulence phenotyping and molecular characterization of a low-pathogenic of *Listeria monocytogenes* from cow's milk. *Acta Biochim. Biophys. Sin.* 38, 262–270.
- Jiang, L., Ke, C., Xu, J., Chen, J., Chen, X., Chen, N., Shuai, J., Fang, W., 2007. *Listeria monocytogenes* mutants carrying Newcastle disease virus F gene fused to its *actA* and *plcB*: in vitro expression and immunogenicity in chickens. *Acta Biochim. Biophys. Sin.* 39, 57–66.
- Jiang, L., Chen, J., Xu, J., Zhang, X., Wang, S., Zhao, H., Vongxay, K., Fang, W., 2008. Virulence characterization and genotypic analyses of *Listeria monocytogenes* isolates from food and processing environments in eastern China. *Int. J. Food Microbiol.* 121, 53–59.
- Kirchner, M., Higgins, D.E., 2008. Inhibition of ROCK activity allows InlF-mediated invasion and increased virulence of *Listeria monocytogenes*. *Mol. Microbiol.* 68, 749–767.
- Liu, D., Ainsworth, A.J., Austin, F.W., Lawrence, M.L., 2003. Characterization of virulent and avirulent *Listeria monocytogenes* strains by PCR amplification of putative transcriptional regulator and internalin genes. *J. Med. Microbiol.* 52, 1065–1070.
- Liu, D., 2004. *Listeria monocytogenes*: comparative interpretation of mouse virulence assay. *FEMS Microbiol. Lett.* 233, 159–164.
- Liu, D., Lawrence, M.L., Wiedmann, M., Gorski, L., Mandrell, R.E., Austin, F.W., Ainsworth, A.J., 2006a. *Listeria monocytogenes* serotype 4b strains belonging to lineages I and III possess distinct molecular features. *J. Clin. Microbiol.* 44, 214–217.
- Liu, D., Lawrence, M.L., Wiedmann, M., Gorski, L., Mandrell, R.E., Ainsworth, A.J., Austin, F.W., 2006b. *Listeria monocytogenes* subgroups IIIA, IIIB and IIIC delineate genetically distinct populations with varied virulence potential. *J. Clin. Microbiol.* 44, 4229–4233.
- Liu, D., Lawrence, M.L., Ainsworth, A.J., Austin, F.W., 2007a. Toward an improved laboratory definition of *Listeria monocytogenes* virulence. *Int. J. Food Microbiol.* 118, 101–115.
- Liu, D., Lawrence, M.L., Ainsworth, A.J., Austin, F.W., 2007b. A multiplex PCR for species- and virulence-specific determination of *Listeria monocytogenes*. *J. Microbiol. Methods* 71, 133–140.
- McLauchlin, J., 1997. The identification of *Listeria* species. *Int. J. Food Microbiol.* 38, 77–81.
- Nelson, K.E., Fouts, D.E., Mongodin, E.F., Ravel, J., DeBoy, R.T., Kolonay, J.F., Rasko, D.A., Angiuoli, S.V., Gill, S.R., Paulsen, I.T., Peterson, J., White, O., Nelson, W.C., Nierman, W., Beanan, M.J., Brinkac, L.M., Daugherty, S.C., Dodson, R.J., Durkin, A.S., Madupu, R., Haft, D.H., Selengut, J., Aken, S.V., Khouri, H., Fedorova, N., Forberger, H., Tran, B., Kathariou, S., Wonderling, L.D., Uhlrich, G.A., Bayles, D.O., Luchansky, J.B., Fraser, C.M., 2004. Whole genome comparisons of serotype 4b and 1/2a strains of the food-borne pathogen *Listeria monocytogenes* reveal new insights into the core genome components of this species. *Nucleic Acids Res.* 32, 2386–2395.
- Olier, M., Pierre, F., Lemaitre, J.P., Divies, C., Rousset, A., Guzzo, J., 2002. Assessment of the pathogenic potential of two *Listeria monocytogenes* human faecal carriage isolates. *Microbiology* 148, 1855–1862.
- Roberts, A., Chan, Y., Wiedmann, M., 2005. Definition of genetically distinct attenuation mechanisms in naturally virulence-attenuated *Listeria monocytogenes* by comparative cell culture and molecular characterization. *Appl. Environ. Microbiol.* 71, 3900–3910.
- Sabet, C., Lecuit, M., Cabanes, D., Cossart, P., Bierne, H., 2005. LPXTG protein InlJ, a newly identified internalin involved in *Listeria monocytogenes* virulence. *Infect. Immun.* 73, 6912–6922.
- Sabet, C., Lecuit, M., Cabanes, D., Cossart, P., Bierne, H., 2008. The *Listeria monocytogenes* virulence factor InlJ is specifically expressed in vivo and behaves as an adhesin. *Infect. Immun.* 76, 1368–1378.
- Schonberg, A., Bannerman, E., Courtieu, A.L., Kiss, R., McLauchlin, J., Shah, S., Whihelms, D., 1996. Serotyping of 80 strains from the WHO multicentre international typing study of *Listeria monocytogenes*. *Int. J. Food Microbiol.* 32, 279–287.
- Suarez, M., Gonzalez-Zorn, B., Vega, Y., Chico-Calero, I., Vazquez-Boland, J., 2001. A role for ActA in epithelial cell invasion by *Listeria monocytogenes*. *Cell. Microbiol.* 3, 853–864.



- Tsai, Y.L., Orsi, R.H., Nightingale, K.K., Wiedmann, M., 2006. *Listeria monocytogenes* internalins are highly diverse and evolved by recombination and positive selection. *Infect. Genet. Evol.* 6, 378–389.
- Vazquez-Boland, J.A., Kuhn, M., Berche, P., Chakraborty, T., Dominguez-Bernal, G., Goebel, W., Gonzalez-Zorn, B., Wehland, J., Kreft, J., 2001. *Listeria* pathogenesis and molecular virulence determinants. *Clin. Microbiol. Rev.* 14, 584–640.
- Wiedmann, M., Bruce, J.L., Keatine, C., Johnson, A.E., McDonough, P.L., Batt, C.A., 1997. Ribotypes and virulence gene polymorphisms suggest three distinct *Listeria monocytogenes* lineages with differences in pathogenic potential. *Infect. Immun.* 65, 2707–2716.
- Zeng, H., Zhang, X., Sun, Z., Fang, W., 2006. Multiplex PCR identification of *Listeria monocytogenes* isolates from milk and milk-processing environments. *J. Sci. Food Agric.* 86, 367–371.